


RHIC Beam Energy Scan Program

Experimental Approach to the QCD Phase Diagram

Grazyna Odyniec / LBNL, Berkeley



central Au+Au @ 7.7 GeV event in STAR TPC



Outline :

Main goal of BES: study QCD phase diagram

Heavy Ion Collisions – the only experimental tool


BES @ RHIC: Physics goals and observables:

- search for the CP and 1st order phase transition
- demonstrate the onset of deconfinement (QGP)

Run 10 – STAR experience

Run 11

USA-NSAC 2007 Long-range Plan






QCD phase diagram - Theory

M. Stephanov, hep-ph/0402115v1 (March 2006)

Theory at the “edges” is believed to be well understood:

1. Lattice QCD finds a smooth crossover at large T and $\mu_B \sim 0$
2. Various models find a strong 1st order transition at large μ_B

So, **there must be a critical point**, but where?



Lattice at $\mu_B \neq 0$: serious problems, several methods on lattice, no agreement so far:

→ CP range: $160 < \mu_B < 500$ MeV


Given the significant theoretical difficulties, data may lead the study of QCD phase diagram

➔ **Beam Energy Scan Program at RHIC will cover this range**



Beam Energy Scan at RHIC: $\sqrt{s_{NN}} \sim 5-50$ GeV

experimental window to QCD phenomenology
at finite temperature and baryon number density



at RHIC : indications of sQGP
but remain unknown:

- boundary between hadronic and partonic phases
- critical point

HOW to investigate it ?

BES @ RHIC


$160 \text{ MeV} < \mu_B < 500 \text{ MeV}$

also: SPS, FAIR (fixed target)



RHIC and BNL from space


RHIC = Relativistic Heavy Ion Collider
Located at BNL= Brookhaven National Laboratory




Landsat-4 satellite in July of 1982

Relativistic Heavy Ion Collider (RHIC)

Brookhaven National Laboratory (BNL), Upton, NY



Relativistic Heavy Ion Collider (RHIC)



- 2 concentric rings of 1740 superconducting magnets
 - 3.8 km circumference
 - counter-rotating beams of ions from p to Au
- max center-of-mass energy: AuAu 200 GeV, pp 500 GeV



BES: Experimental Program

<http://drupal.star.bnl.gov/STAR/starnotes/public/sn0493>

Search for:

(1) indications of the existence of Critical Point & phase transition

- fluctuation measures
 - higher moments of net proton distribution (kurtosis) ★
- azimuthally-sensitive femtoscopy
- elliptic & directed flow
- ...

(2) disappearance of signals of partonic degrees of freedom seen at 200 GeV



- disappearance of constituent-quark-number scaling of v_2 ★
- disappearance of hadron suppression in central collisions
- disappearance of ridge
- local parity violation
- ...



Critical Point search – Fluctuations maximized at CP

example: e-by-e fluctuations in K/ π ratio

PRL 103, 092301 (2009)




NA49 fluct. > STAR fluct. ?

CP at lower energies ? (but diff. acceptance).

Grazyna Odyniec

more sensitive : - Higher Moments



Thermodynamics: Divergence of susceptibilities for conserved quantities (B,Q,S) at critical point.

Lattice QCD: Spikes for both χ_B and χ_S

Berdnikov, Rajagopal, PRD61, 105017 (00)

Stephanov, Rajagopal, Shuryak, PRD 60, 114028 (99)


Hatta, Stephanov, PRL. 91, 102003 (03)

Gavai and Gupta, Phys. Rev. D 78, 114503 (2008);

Gupta, arXiv:0909.4630 [nucl-ex].

$$\chi_2^X = \frac{1}{VT^3} \langle N_X^2 \rangle$$

$$\chi_4^X = \frac{1}{VT^3} (\langle N_X^4 \rangle - 3\langle N_X^2 \rangle^2)$$



Observable:

Kurtosis of net-proton & net-C



- connect to lattice calculations!
- sensitive to long range fluctuations

Caveats: dynamical effects in collisions

- finite time and size
- critical slowing

High Moment Analysis (BES)

STAR: aXiv:1004.4959, sub. to PRL



High moments are more sensitive to critical point related fluctuation.



The 4th moment, Kurtosis, is directly related to the corresponding thermodynamic quantity: susceptibility for conserved quantum numbers such as Baryon number, charge, strangeness...



Centrality dependence of net-proton Kurtosis

STAR Preliminary:

200 GeV Au+Au ($\mu_B \leq 25$ MeV)



First Kurtosis measurement for net-protons in high-energy nuclear collisions

Monotonic behavior observed at relatively small μ_B region \rightarrow baseline

Grazyna Odyniec




Disappearance of partonic degrees of freedom (I)

(Onset of sQGP)

disappearance of n_q scaling, disappearance of hadron suppression at high p_T , ... (a long list)


n_q scaling observed at RHIC:




- (1) Mass separation at low p_T
- (2) Light and heavy quarks have similar magnitude of flow
- (3) In intermediate p_T : separation between baryon and meson band



Disappearance of partonic degrees of freedom (II)



Scaling flow parameters by quark content n_q (baryons=3, mesons=2) resolves meson-baryon separation of final state hadrons




flow developed in pre-hadronic stage
DECONFINEMENT at RHIC

With lowering energy, disappearance of n_q scaling would suggest that we exit partonic dof world



Will we be able to see it ?


PRL 92, 052302(04), 95, 122301(05), nucl-ex/0405022, QM05



Yes, a few M is enough !

Local Parity Violations in Deconfined Medium

D.E. Kharzeev et al, NPA 803, 227 (2008)
 K. Fukushima et al, PRD 78, 074033 (2008)




- (1) Under strong magnetic field, when the system is in the state of **deconfinement** and **chiral symmetry restoration** is reached, local fluctuation may lead to parity violation.
- (2) Experimentally one would observe the separation of the charges in high-energy nuclear collisions.

Parity even observable: $\langle \cos(\phi_\alpha + \phi_\beta - 2\Psi_{RP}) \rangle$
 Voloshin, PR C62, 044901(00),

- (3) In RHIC Beam Energy Scan program - test the model prediction

- the energy when the charge separation disappear => phase boundary





Collision Energies (GeV)	Millions of Events Needed					
	5	7.7	11.5	17.3	27	39
Observables	Millions of Events Needed					
v_2 (up to ~ 1.5 GeV/c)	0.3	0.2	0.1	0.1	0.1	0.1
v_1	0.5	0.5	0.5	0.5	0.5	0.5
Azimuthally sensitive HBT	4	4	3.5	3.5	3	3
PID fluctuations (K/ π)	1	1	1	1	1	1
net-proton kurtosis	5	5	5	5	5	5
differential corr & fluct vs. centrality	4	5	5	5	5	5
n_q scaling $\pi/K/p/\Lambda$ ($m_T - m_0$)/ $n < 2$ GeV	8.5	6	5	5	4.5	4.5
ϕ/Ω up to $p_T/n_q = 2$ GeV/c		56	25	18	13	12
R_{CP} up to $p_T \sim 4.5$ GeV/c (at 17.3) 5.5 (at 27) & 6 GeV/c (at 39)				15	33	24
untriggered ridge correlations		27	13	8	6	6
parity violation		5	5	5	5	5



<http://www.bnl.gov/npp/pac.asp>

Recommendations of BNL Nuclear and Particle Physics Program Advisory Committee (PAC):

Run 10 (2010):


1. 10 weeks of Au+Au at 200 GeV
2. 12 weeks for a beam energy scan (BES) with Au+Au collisions:
 1. - 4 weeks 62 GeV
 2. - 8 weeks lower energies
 1. 0.5 week 39 and 27 GeV
 2. 1 week at 18 GeV (10 M)
 3. 2 weeks at 11 GeV (6 M)
 4. 4 weeks at 7.7 GeV (3.6 M)

Sufficient rates for the initial physics program at all energies

“binary” experiment: YES/NO (no “maybe’s” & more statistics needed)





STAR experience with low energy running





STAR experiment demonstrated capabilities



only a few 10^3 events taken during machine test



9.2 GeV results consistent with the published data

STAR : PRC 79 (2009) 034909, arXiv: 0903.4702

NA49 : PRC 66 (2002) 054902, PRC 77 (2008) 024903, PRC 73 (2006) 044910

E802(AGS) : PRC 58 (1998) 3523, PRC 60 (1999) 044904, PRC 62 (2000) 024901, PRC 68 (2003) 054903


Elliptic Flow



STAR and NA49 results are consistent
 STAR 9.2GeV v_2 fits with the observed trends

NA49 : PRC 68 (2003) 034903
 AGS : PLB 474 (2000) 27
 STAR : PRC 77 (2008) 054901 : PRC 75 (2007)
 054906, PRC 72 (2005) 014904
 PHOBOS : PRC 72 (2005) 051901 :
 PRL 98 (2007) 242302
 PHENIX : PRL 98 (2007) 162301

Pion Interferometry



π^-

error bars for Au+Au 9.2 GeV are statistical
systematic errors < 10 % for all radii

STAR : PRC 71 (2005) 044906, PRL 87 (2001) 082301

PHENIX : PRL 88 (2002) 192302, PRL 93(2004) 152302

E802 : PRC 66 (2002) 054906 NA44 : PRC 58 (1998) 1656

CERES : NPA 714 (2003) 124 E866 : NPA 661 (1999) 439



E895 : PRL 84 (2000) 2798 NA49 : PRC 77 (2008) 64908

PHOBOS : PRC 73 (2006) 031901 WA97 : JPG 27 (2001) 2325

Phys. Rev. C 81, 024911(2010)

STAR was ready for run 10 !
to take up Beam Energy Scan Program

STAR Experimental Configuration for Run 10 (2010)



No SVT (low material)
 100% TOF
 FTPCs
 Large beam pipe



Run 10 – part I of BES@RHIC

Hardware and operation improvements

Main directions of Beam Energy Scan program at RHIC established:

- search for turn-off of sQGP signatures
- search for the evidence of CP and/or 1st order phase transition
- + many other measurements

Strategy: scan available phase space with (6) equally spaced points between 5 and 39 GeV (we already have 62, 130, 200 data), and return to “interesting” regions for more detailed studies in the next year



Train left the station on April 8th
with 39 GeV Au+Au collisions ...


Run 10 :

39, 7.7 and 11.5 GeV Au+Au


to be continued (run 11) next year




Central Au+Au @ 7.7 GeV Event



G




Typical Au+Beam-
pipe @ 3.85 GeV
event



Event outside active
TPC volume




Au+Au @ 7.7 AGeV - vertex reconstruction – bck !



- Au+beam-pipe events from the beam halo



Au+Au @ 7.7 AGeV – sources of background







Two flanges at ± 4.4 m outside the TPC produced most of the background at the first days

Grazyna Odyniec

vertex distributions though the run ...

middle: Day 130

end: Day 147





Grazyna

Superb improvement !




High Level Trigger (HLT): Vertex



HLT is able to reconstruct online the primary vertices
HLT good event rate is very close to offline QA rate
Priceless redundancy !






Online HLT good event rate



~ 9 Hz !



STAR data summary AuAu @ 7.7 GeV



Collected 5.014 M events

– met the goal !

Operated in 10 minutes stores environment

Average uptime ~ 11 hours

Collecting up to 350 k events per day

Grazyna Odyniec



Statistics from Run 10




Beam Energy ($\sqrt{s_{NN}}$, GeV)	Minbias (Million)	Central (Million)	High-Tower Sampled Luminosity	FTPC+PMD (Million)
200	355	265	2.6 (nb ⁻¹)	5
62.4	140	33	175 (μb ⁻¹)	3.5
39	250		62 (μb ⁻¹)	23
7.7	5	N/A	N/A	N/A
11.5	≥ 7.5	N/A	N/A	N/A
5	Commissioning	N/A	N/A	N/A

Identified Particle Acceptance at STAR


Au+Au at 7.7 GeV

Au+Au at 39 GeV

Au+Au at 200 GeV




Multiplicity at 7.7, 39, and ... GeV






STAR Performance in Run 10 Particle Identification at 7.7 GeV

TPC PID








TPC+TOF PID




Particle Identification – part II

$\sqrt{s_{NN}} = 39 \text{ GeV Au + Au Collisions}$



Invariant Mass (GeV)

Event Plane Resolutions



Event plan measurements at STAR:

- (1) TPC ($|\eta| \leq 1$)
- (2) FTPC ($2.5 \leq |\eta| \leq 4.2$)
- (3) BBC ($3.8 \leq |\eta| \leq 5.2$)



Summary

- RHIC Beam Energy Scan - **Fantastic success !**
Au + Au at 39, 7.7 and 11.5 GeV runs:
 - Met all goals and far exceeded for some data points
 - 7.7 GeV (34 days) and 11.5 GeV (8+3 days) : $N_{\text{events}} > 5 \text{ M}$
 - 39 GeV (15 days): $N_{\text{events}} \sim 170 \text{ M events}$
 - Dramatic improvement of the collider performance at 7.7 GeV
- Preliminary results based on fast offline run 10 data look very promising
- Final calibration results soon
- **Last call for predictions on critical point !!!**
- PAC in two weeks ... discussion of run 11 begins !



STAR BES for Runs 11 and 12

Run	Beam Energy	Time	System	Goal	New Detector
11	$\sqrt{s_{NN}} = 18, 27 \text{ GeV}$	2 weeks	Au + Au	100, 150M minbias	HLT
	$\sqrt{s_{NN}} = 200 \text{ GeV}$	4 weeks	U + U	200M minbias 200M central	
	$\sqrt{s} = 500 \text{ GeV}$	5 weeks	$p_{\uparrow} p_{\uparrow}$	trans. $P^2 * L = 4 \text{ pb}^{-1}$	
		6 weeks	$p_{\rightarrow} p_{\rightarrow}$	long. $P^2 * L = 20 \text{ pb}^{-1}$	
	1 week	$p_{\uparrow} p_{\uparrow}$	pp2pp at high β^*		
12	$\sqrt{s} = 500 \text{ GeV}$	10 weeks	$p_{\rightarrow} p_{\rightarrow}$	long. $P^2 * L = 50 \text{ pb}^{-1}$ $P^4 * L = 15 \text{ pb}^{-1}$	FGT
	or		or	or	
	$\sqrt{s} = 200 \text{ GeV}$		$p_{\uparrow} p_{\uparrow}$ $p_{\rightarrow} p_{\rightarrow}$	trans. $P^2 * L = 8.5 \text{ pb}^{-1}$ long. $P^4 * L = 4.3 \text{ pb}^{-1}$	
	$\sqrt{s_{NN}} = 200 \text{ GeV}$		U + U or Au+Au	3.5 nb^{-1} U+U or 5 nb^{-1} Au+Au	MTD



Thank you!

